

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

ADAM15 RABBIT PAB

货号: S216938

产品全名: ADAMI5 兔多抗

基因符号 MDC15

UNIPROT ID: Q13444 (Gene Accession - BC014566)

背景: The protein encoded by this gene is a member of the ADAM (a disintegrin and metalloproteinase) protein family. ADAM family members are type I transmembrane glycoproteins known to be involved in cell adhesion and proteolytic ectodomain processing of cytokines and adhesion molecules. This protein contains multiple functional domains including a zinc-binding metalloprotease domain, a disintegrin-like domain, as well as a EGF-like domain. Through its disintegrin-like domain, this protein specifically interacts with the integrin beta chain, beta 3. It also interacts with Src family protein-tyrosine kinases in a phosphorylation-dependent manner, suggesting that this protein may function in cellcell adhesion as well as in cellular signaling. Multiple alternatively spliced transcript variants encoding distinct isoforms have been observed.

抗原: Fusion protein of human ADAM15

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-200;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit


克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat


成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Signal Transduction, Cell Biology


Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human tonsil In comparision with the IHC on the left, the same paraffin-embedded tissue using 216938(ADAM15 Antibody) at a dilution of 1/70(Cytoplasm).



Human tonsil tissue is first treated with the fusion protein and then with 216938 (Anti-ADAM15 Antibody) at dilution 1/70.



Gel: 6%SDS-PAGE, Lysate: 40 µg;

Lane 1-5: NIH/3T3, HepG2, PC-3, Jurkat cell, Mouse heart tissue

Primary antibody: 216938(ADAM15 Antibody) at dilution 1/800; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000

dilution:

Exposure time: 5 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010