

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **AKAP8L RABBIT PAB**


货号: S216964 产品全名: AKAP8L 兔多抗 基因符号 HA95; HAP95; NAKAP; NAKAP95 UNIPROT ID: Q9ULX6 (Gene Accession - BC000713) 背景: A-kinase anchor protein 8-like is an enzyme that in humans is encoded by the AKAP8L gene. Could play a role in constitutive transport element (CTE)-mediated gene expression. Does not seem to be implicated in the binding of regulatory subunit II of PKA. May be involved in nuclear envelope breakdown and chromatin condensation. May regulate the initiation phase of DNA replication when associated with TMPObeta. 抗原: Fusion protein of human AKAP8L 经过测试的应用: ELISA, WB, IHC 推荐稀释比: IHC: 25-100;WB: 200-1000;ELISA: 2000-5000 种属反应性: Rabbit 克隆性: Rabbit Polyclonal 亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol 研究领域: Epigenetics and Nuclear Signaling 储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human breast cancer tissue using 216964(AKAP8L Antibody) at a dilution of 1/15(Nucleus).




The image on the left is immunohistochemistry of paraffinembedded Human ovarian cancer tissue using 216964(Anti-AKAP8L Antibody) at a dilution of 1/15.



In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with the fusion protein and then with 216964(Anti-AKAP8L Antibody) at dilution 1/15.



In comparision with the IHC on the left, the same paraffin-embedded Human ovarian cancer tissue is first treated with fusion protein and then with D221552(Anti-AKAP8L Antibody) at dilution 1/15.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-2: Jurkat cells, 293T cells; Primary antibody: 216964(AKAP8L Antibody) at dilution 1/84; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 20 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010