

## AMIGO2 RABBIT PAB

货号: S216996

产品全名: AMIGO2 兔多抗

基因符号: ALII; DEGA; AMIGO-2

**UNIPROT ID:** Q86SJ2 (Gene Accession - BC047595 )

**背景:** Predicted to be involved in several processes, including heterophilic cell-cell adhesion via plasma membrane cell adhesion molecules; homophilic cell adhesion via plasma membrane adhesion molecules; and negative regulation of programmed cell death. Predicted to act upstream of or within positive regulation of synapse assembly. Predicted to be located in nucleus and plasma membrane. Predicted to be integral component of membrane. Biomarker of gastric adenocarcinoma.

**抗原:** Fusion protein of human AMIGO2

**经过测试的应用:** ELISA, WB, IHC

**推荐稀释比:** IHC: 50-200;WB: 500-2000;ELISA: 5000-10000

**种属反应性:** Rabbit

**克隆性:** Rabbit Polyclonal

**亚型:** Immunogen-specific rabbit IgG


**纯化:** Antigen affinity purification

**种属反应性:** Human, Mouse, Rat


**成分:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

**研究领域:** Neuroscience


**储存和运输:** Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 216996 (AMIGO2 Antibody) at a dilution of 1/60 (Cytoplasm).



In comparison with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the fusion protein and then with 216996 (Anti-AMIGO2 Antibody) at dilution 1/60.



Gel: 8%SDS-PAGE, Lysate: 40 µg;

Lane: SP20 cell lysate;

Primary antibody: 216996 (AMIGO2 Antibody) at dilution 1/800;

Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;

Exposure time: 90 seconds