

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010


## **ARFRP1 RABBIT PAB**

货号: S214084 产品全名: ARFRPI 兔多抗 基因符号 ARP; Arp1; ARL18 UNIPROT ID: Q13795 (Gene Accession - NP\_003215) 背景: The protein encoded by this gene is a membrane-associated GTP-ase which localizes to the plasma membrane and is related to the ADP-ribosylation factor (ARF) and ARF-like (ARL) proteins. This gene plays a role in membrane trafficking between the trans-Golgi network and endosomes. Alternatively spliced transcript variants encoding different isoforms have been identified. 抗原: Synthetic peptide of human ARFRP1 经过测试的应用: ELISA, WB, IHC 推荐稀释比: IHC: 25-100;WB: 500-2000;ELISA: 2000-5000 种属反应性: Rabbit 克隆性: Rabbit Polyclonal 亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat 成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol 研究领域: Signal Transduction 储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human breast cancer tissue using 214084(ARFRP1 Antibody) at a dilution of 1/25(Cytoplasm).



The image on the left is immunohistochemistry of paraffinembedded Human cervical cancer tissue using 214084(Anti-ARFRP1 Antibody) at a dilution of 1/25. In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with the synthetic peptide and then with 214084(Anti-ARFRP1 Antibody) at dilution 1/25.



In comparision with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with synthetic peptide and then with D161316(Anti-ARFRP1 Antibody) at dilution 1/25.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-4: 293T cells, human fetal liver tissue, human cervical cancer tissue, Human fetal muscle tissue; Primary antibody: 214084(ARFRP1 Antibody) at dilution 1/800; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 40 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010