

## BID RABBIT PAB

货号: S216370

产品全名: BID 兔多抗

基因符号 FP497

**UNIPROT ID:** P55957 (Gene Accession - BC022072 )

**背景:** This gene encodes a death agonist that heterodimerizes with either agonist BAX or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH-terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but the full-length nature of some variants has not been defined.

**抗原:** Fusion protein of human BID

**经过测试的应用:** ELISA, WB, IHC

**推荐稀释比:** IHC: 25-100;WB: 200-2000;ELISA: 1000-5000

**种属反应性:** Rabbit

**克隆性:** Rabbit Polyclonal

**亚型:** Immunogen-specific rabbit IgG



**纯化:** Antigen affinity purification

**种属反应性:** Human

**成分:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


**研究领域:** Cancer

**储存和运输:** Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human renal cancer tissue using 216370 (BID Antibody) at a dilution of 1/25 (Cytoplasm, Cell membrane).

In comparison with the IHC on the left, the same paraffin-embedded Human renal cancer tissue is first treated with the fusion protein and then with 216370 (Anti-BID Antibody) at dilution 1/25.



Gel: 12% SDS-PAGE, Lysate: 30 µg;

Lane 1-3: Human fetal brain tissue, HT29 cells, HepG2 cells;

Primary antibody: 216370 (BID Antibody) at dilution 1/220;

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;

Exposure time: 10 seconds