

BIRC3 RABBIT PAB

货号: S216817

产品全名: BIRC3 兔多抗

基因符号: AIP1; AIP2; MIHC; CIAP2; HAIPI; HIAP1; IAP-1; MALT2; RNF49; c-IAP2

UNIPROT ID: Q13489 (Gene Accession - BC037420)

背景: This gene encodes a member of the IAP family of proteins that inhibit apoptosis by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. It contains 3 baculovirus IAP repeats and a ring finger domain. Transcript variants encoding the same isoform have been identified.

抗原: Fusion protein of human BIRC3

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-100;WB: 200-1000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Cancer


储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 216817(BIRC3 Antibody) at a dilution of 1/75(Cytoplasm or Nucleus).



In comparison with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the fusion protein and then with 216817(Anti-BIRC3 Antibody) at dilution 1/75.




The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using 216817(Anti-BIRC3 Antibody) at a dilution of 1/75.



In comparison with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with fusion protein and then with D221291(Anti-BIRC3 Antibody) at dilution 1/75.

kDa



Gel: 8%SDS-PAGE, Lysate: 40 µg;
Lane: Raji cell lysate;
Primary antibody: 216817(BIRC3 Antibody) at dilution 1/250;
Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;
Exposure time: 2 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010
