

BTG2 RABBIT PAB

货号: S216378

产品全名: BTG2 兔多抗

基因符号: PC3; APRO1; TIS21

UNIPROT ID: P78543 (Gene Accession - BC105949)

背景: The protein encoded by this gene is a member of the BTG/Tob family. This family has structurally related proteins that appear to have antiproliferative properties. This encoded protein is involved in the regulation of the G1/S transition of the cell cycle.

抗原: Fusion protein of human BTG2

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 100-300;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human, Mouse, Rat


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Epigenetics and Nuclear Signaling, Neuroscience


储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human cervical cancer tissue using 216378(BTG2 Antibody) at a dilution of 1/95(Cytoplasm).



In comparison with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with the fusion protein and then with 216378(Anti-BTG2 Antibody) at dilution 1/95.



Gel: 12%SDS-PAGE, Lysate: 40 µg;
Lane 1-7: Mouse lung tissue, Jurkat, 293T, K562 cell, Mouse brain tissue, Mouse kidney tissue, 231 cell lysates;
Primary antibody: 216378(BTG2 Antibody) at dilution 1/500;
Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;
Exposure time: 1 minute