

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **C8G RABBIT PAB**

货号: S218113 产品全名: C8G 兔多抗 基因符号 C8C UNIPROT ID: P07360 (Gene Accession - BC113626) 背景: The protein encoded by this gene belongs to the lipocalin family. It is one of the three subunits that constitutes complement component 8 (C8) which is compared of a disulfide-linked C8 alpha-agamma bateradimer and a pop-covalently associated C8 b

component 8 (C8), which is composed of a disulfide-linked C8 alpha-gamma heterodimer and a non-covalently associated C8 beta chain. C8 participates in the formation of the membrane attack complex (MAC) on bacterial cell membranes. While subunits alpha and beta play a role in complement-mediated bacterial killing, the gamma subunit is not required for the bactericidal activity. 抗原: Full length fusion 蛋白

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 200-1000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG

纯化: Antigen affinity purification

种属反应性: Human

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Immunology

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010



Immunchistochemistry analysis of paraffin embedded Human cervical cancer tissue using 218113(C8G Antibody) at a dilution of 1/25(Cytoplasm).




In comparision with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with the fusion protein and then with 218113(Anti-C8G Antibody) at dilution 1/25.



The image on the left is immunohistochemistry of paraffinembedded Human tonsil tissue using 218113(Anti-C8G Antibody) at a dilution of 1/25.

In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with fusion protein and then with D223748(Anti-C8G Antibody) at dilution 1/25.



Gel: 12%SDS-PAGE, Lysate: 40 µg; Lane: Human colorectal cancer tissue; Primary antibody: 218113(C8G Antibody) at dilution 1/200; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 10 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010