

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **CA7 RABBIT PAB**

货号: S218418 产品全名: CA7 兔多抗 基因符号 CAVII

UNIPROT ID: P43166 (Gene Accession - BC033865)

背景: Carbonic anhydrases are a large family of zinc metalloenzymes that catalyze the reversible hydration of carbon dioxide. They participate in a variety of biological processes, including respiration, calcification, acid-base balance, bone resorption, and the formation of aqueous humor, cerebrospinal fluid, saliva, and gastric acid. They show extensive diversity in tissue distribution and in their subcellular localization. The cytosolic protein encoded by this gene is predominantly expressed in the salivary glands. Alternative splicing in the coding region results in multiple transcript variants encoding different isoforms.

抗原: Full length fusion 蛋白 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-300;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit 克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Cell Biology

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human gastric cancer tissue using 218418(CA7 Antibody) at a dilution of 1/50(Cytoplasm or Nucleus).




In comparision with the IHC on the left, the same paraffin-embedded Human gastric cancer tissue is first treated with the fusion protein and then with 218418(Anti-CA7 Antibody) at dilution 1/50.



The image on the left is immunohistochemistry of paraffinembedded Human thyroid cancer tissue using 218418(Anti-CA7 Antibody) at a dilution of 1/50.



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with fusion protein and then with D224370(Anti-CA7 Antibody) at dilution 1/50.



Gel: 12%SDS-PAGE, Lysate: 40 µg; Lane 1-4: Human urinary bladder tissue, Mouse heart tissue, Mouse kidney tissue and RAW264.7 cell lysates; Primary antibody: 218418(CA7 Antibody) at dilution 1/500; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 5 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010