

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## CACNA1G RABBIT PAB

货号: S220417 产品全名: CACNAIG 兔多抗 基因符号 NBR13; Cav3.1; Ca(V)T.1

UNIPROT ID: 043497 (Gene Accession - NP\_061496)

背景: Voltage-dependent calcium channels mediate the entry of calcium ions into excitable cells, and are also involved in a variety of calcium-dependent processes, including muscle contraction, hormone or neurotransmitter release, and gene expression. Calcium channels are multisubunit complexes composed of alpha-1, beta, alpha-2/delta, and gamma subunits. The channel activity is directed by the pore-forming alpha-1 subunit, whereas, the others act as auxiliary subunits regulating this activity. The distinctive properties of the calcium channel types are related primarily to the expression of a variety of alpha-1 isoforms, alpha-1A, B, C, D, E, and S. This gene encodes the alpha-1A subunit, which is predominantly expressed in neuronal tissue. Mutations in this gene are associated with 2 neurologic disorders, familial hemiplegic migraine and episodic ataxia 2.

抗原: Synthetic peptide of human CACNAIG

经过测试的应用: ELISA, IHC

推荐稀释比: IHC: 15-50; ELISA: 1000-2000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG

纯化: Antigen affinity purification

种属反应性: Human, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 220417(CACNA1G Antibody) at a dilution of 1/20(Cytoplasm).



The image on the left is immunohistochemistry of paraffinembedded Human liver cancer tissue using 220417(Anti-CACNAIG Antibody) at a dilution of 1/20.



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the synthetic peptide and then with 220417(Anti-CACNA1G Antibody) at dilution 1/20.



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with synthetic peptide and then with D261501(Anti-CACNA1G Antibody) at dilution 1/20.