

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **CALCOCO1 RABBIT PAB**

货号: S218907

产品全名: CALCOCOI 兔多抗

基因符号 Cocoa; PP13275; calphoglin

UNIPROT ID: Q9P1Z2 (Gene Accession - BC003177)

背景: CALCOCOI (calcium-binding and coiled-coil domain-containing protein I), also known as cocoa, calphoglin, sarcoma antigen NY-SAR-3 or coiled-coil coactivator protein, is a 691 amino acid protein that shuttles between the cytoplasm and nucleus and functions as coactivator for aryl hydrocarbon and nuclear receptors. A member of the CALCOCO family, CALCOCOI is forms a calphoglin complex with PPAI and PGM I and contains multiple functional domains through which it acts as a component of both the androgen signaling pathway and the Wnt/?-catenin signaling pathway. CALCOCOI exists as three alternatively spliced isoforms (termed Q9PIZ2-1, 2 and 3), which are encoded by genes mapping to human chromosome 12q13.13 and mouse chromosome 15 F3.

抗原: Fusion protein of human CALCOCOI

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 40-200;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction, Epigenetics and Nuclear Signaling 储存和运输: Store at -20°C. Avoid repeated freezing and thawing



## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010






Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 218907(CALCOCOI Antibody) at a dilution of 1/50(Cytoplasm or Nucleus).


In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the fusion protein and then with 218907(Anti-CALCOCOI Antibody) at dilution 1/50.



The image on the left is immunohistochemistry of paraffinembedded Human esophagus cancer tissue úsing 218907(Anti-CALCOCOI Antibody) at a dilution of 1/50.



In comparision with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with fusion protein and then with D225479(Anti-CALCOCOI Antibody) at dilution 1/50.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-5: 231, Jurkat, Hela, 293T cell and Mouse heart tissue lysates; Primary antibody: 218907(CALCOCOI Antibody) at dilution 1/300; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 5 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010