

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

CASP10 RABBIT PAB

货号: S219607 产品全名: CASP10 兔多抗 基因符号 MCH4; ALPS2; FLICE2; FLICE-2 UNIPROT ID: Q92851 (Gene Accession - NP_116759)

背景: This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolyti

caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 3 and 7, and the protein itself is processed by caspase 8. Mutations in this gene are associated with type IIA autoimmune lymphoproliferative syndrome, non-Hodgkin lymphoma and gastric cancer. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.

抗原: Synthetic peptide of human CASP10

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-200;WB: 200-1000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification


种属反应性: Human

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Cancer, Cell Biology


储存和运输: Store at -20°C. Avoid repeated freezing and thawing





Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 219607(CASP10 Antibody) at a dilution of 1/50(Cytoplasm).

In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the synthetic peptide and then with 219607(Anti-CASP10 Antibody) at dilution 1/50.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: HepG2 cell lysate; Primary antibody: 219607(CASP10 Antibody) at dilution 1/550; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution; Exposure time: 5 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010