

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

CCNE1 RABBIT PAB

货号: S219732

产品全名: CCNEI 兔多抗 基因符号 CCNE; pCCNEI

UNIPROT ID: P24864 (Gene Accession - NP_001229)

背景: The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle GI/S transition. This protein accumulates at the GI-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB.

抗原: Synthetic peptide of human CCNEI

经过测试的应用: ELISA, IHC

推荐稀释比: IHC: 30-150; ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification

种属反应性: Human

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Epigenetics and Nuclear Signaling, Cancer

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 219732(CCNE1 Antibody) at a dilution of 1/25(Nucleus).



In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the synthetic peptide and then with 219732(Anti-CCNEI Antibody) at dilution 1/25.



The image on the left is immunohistochemistry of paraffinembedded Human cervical cancer tissue using 219732(Anti-CCNEI Antibody) at a dilution of 1/25.



In comparision with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with synthetic peptide and then with D260239(Anti-CCNEI Antibody) at dilution 1/25.