CCNE2 RABBIT PAB

货号: S213471
产品全名: CCNE2 兔多抗
基因符号 CYCE2
UNIPROT ID: O96020 (Gene Accession – NP_477097)
背景: The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2. This cyclin has been shown to specifically interact with CIP/KIP family of CDK inhibitors, and plays a role in cell cycle G1/S transition.
抗原: Synthetic peptide of human CCNE2
经过测试的应用: ELISA, IHC
推荐稀释比: IHC: 50-200; ELISA: 2000-5000
种属反应性: Rabbit
克隆性： Rabbit Polyclonal
亚型: Immunogen-specific rabbit IgG
纯化: Antigen affinity purification
种属反应性: Human
成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol
研究领域: Epigenetics and Nuclear Signaling, Cancer
储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human esophagus cancer tissue using 213471(CCNE2 Antibody) at a dilution of 1/25(Nucleus).



In comparision with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with the synthetic peptide and then with 213471(Anti-CCNE2 Antibody) at dilution 1/25.



The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using 213471(Anti-CCNE2 Antibody) at a dilution of 1/25.



In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with synthetic peptide and then with D160240(Anti-CCNE2 Antibody) at dilution 1/25.

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC APPLICATIONS

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC APPLICATIONS