

## CD22 RABBIT MAB

货号: N261986

产品全名: CD22 兔单克隆抗体

基因符号 CD22; SIGLEC2; B-cell receptor CD22; B-lymphocyte cell adhesion molecule; BL-CAM; Sialic acid-binding Ig-like lectin 2; Siglec-2; T-cell surface antigen Leu-14; CD antigen CD22

**UNIPROT ID:** P20273

**背景:** Acts as a regulator of B cell signaling. CD22 is expressed as both a cytoplasmic and membrane protein during discrete stages of B cell lymphocyte differentiation. The cytoplasmic form of CD22, expressed early in B cell development, is a useful marker for acute lymphocytic leukemia. The membrane form of CD22 is expressed in mature B cells prior to their differentiation into plasma cells. Alternative splicing results in two different isoforms, CD22 $\alpha$  and CD22 $\beta$ .

**抗原:** Recombinant protein of human CD22

**经过测试的应用:** WB, ICC/IF, IP

**推荐稀释比:** WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/20

**种属反应性:** Rabbit

**克隆性:** Rabbit Monoclonal

**克隆编号:** R01-8D2

**分子量:** Calculated MW: 95 kDa; Observed MW: 140 kDa

**亚型:** IgG


**纯化:** Affinity Purified

**种属反应性:** Human


**成分:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

**研究领域:** Immunology

**储存和运输:** Store at -20°C. Avoid repeated freezing and thawing



Western blot analysis of CD22 in Raji, K562, Ramos lysates using CD22 antibody.



Immunocytochemistry analysis of CD22 (green) in HL-60 using CD22 antibody, and DAPI (blue).