

CD62P RABBIT PAB

货号: N225044

产品全名: CD62P 兔多抗

基因符号 SELP; GMRP; GRMP; P-selectin; CD62 antigen-like family member P; Granule membrane protein 140; GMP-140; Leukocyte-endothelial cell adhesion molecule 3; LECAM3; Platelet activation dependent granule-external membrane protein; PADGEM; CD62P

UNIPROT ID: P16109

背景: Ca(2+)-dependent receptor for myeloid cells that binds to carbohydrates on neutrophils and monocytes. Mediates the interaction of activated endothelial cells or platelets with leukocytes. The ligand recognized is sialyl-Lewis X. Mediates rapid rolling of leukocyte rolling over vascular surfaces during the initial steps in inflammation through interaction with PSGL1.

抗原: The antiserum was produced against synthesized peptide derived from the N-terminal region of human SELP. AA range:81-130

经过测试的应用: WB,IHC-P,ELISA

推荐稀释比: WB: 1/500-1/1000 IHC: 1/50-1/100 ELISA: 1/10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

分子量: Calculated MW: 91 kDa; Observed MW: 91 kDa

亚型: IgG


纯化: Affinity Purified

种属反应性: Human, Mouse and Rat


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

研究领域: Cardiovascular


储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin-embedded Human tonsils using CD62P antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of CD62P in NIH3T3, KB lysates using CD62P antibody.



Western blot analysis of CD62P in 3T3 lysates using P-Selectin antibody.



Immunohistochemistry analysis of paraffin-embedded Human tonsils using CD62P antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.