

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

CD63 RABBIT PAB

货号: S220112 产品全名: CD63 免多抗 基因符号 MLA1; ME491; LAMP-3; OMA81H; TSPAN30

UNIPROT ID: P08962 (Gene Accession - NP_001244318)

背景: The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. The encoded protein is a cell surface glycoprotein that is known to complex with integrins. It may function as a blood platelet activation marker. Deficiency of this protein is associated with Hermansky-Pudlak syndrome. Also this gene has been associated with tumor progression. Alternative splicing results in multiple transcript variants encoding different protein isoforms.

抗原: Synthetic peptide of human CD63

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-200;WB: 1000-5000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification


种属反应性: Human

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Cardiovascular, Immunology, Cell Markers, Cancer

储存和运输: Store at -20°C. Avoid repeated freezing and thawing





Immunohistochemistry analysis of paraffin embedded Human breast cancer tissue using 220112(CD63 Antibody) at a dilution of 1/50(Cytoplasm).

In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with the synthetic peptide and then with 220112(Anti-CD63 Antibody) at dilution 1/50.

., = = (=) == = = = =	
kDa	
100 —	
70 —	
55—	
35 —	
25—	
15—	

10-

Gel: 12%SDS-PAGE, Lysate: 40 µg; Lane: LNCAP cell lysate; Primary antibody: 220112(CD63 Antibody) at dilution 1/1000; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution; Exposure time: 20 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010