CDK5RAP3 RABBIT PAB

货号: S217272
产品全名: CDK5RAP3 兔多抗
基因符号 C53; IC53; LZAP; HSF-27; MST016; PP1553; OK/SW-cl.114
UNIPROT ID: Q96JB5 (Gene Accession – BC009957)
背景: This gene encodes a protein that has been reported to function in signaling pathways governing transcriptional regulation and cell cycle progression. It may play a role in tumorigenesis and metastasis. A pseudogene of this gene is located on the long arm of chromosome 20. Alternative splicing results in multiple transcript variants that encode different isoforms.
抗原: Fusion protein of human CDK5RAP3
经过测试的应用: ELISA, WB, IHC
推荐稀释比: IHC: 25-100;WB: 500-2000;ELISA: 2000-5000
种属反应性: Rabbit
克隆性： Rabbit Polyclonal
亚型: Immunogen-specific rabbit IgG
纯化: Antigen affinity purification
种属反应性: Human, Mouse, Rat
成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol
研究领域: Signal Transduction, Cancer
储存和运输: Store at -20°C. Avoid repeated freezing and thawing

Product Description

Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 217272(CDK5RAP3 Antibody) at a dilution of 1/40(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the fusion protein and then with 217272(Anti-CDK5RAP3 Antibody) at dilution 1/40.



The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using 217272(Anti-CDK5RAP3 Antibody) at a dilution of 1/40.



In comparision with the IHC on the left, the same paraffin-embedded Human colon cancer tissue is first treated with fusion protein and then with D222083(Anti-CDK5RAP3 Antibody) at dilution 1/40.



Gel: 6%SDS-PAGE, Lysate: 40 µg;
Lane 1-2: Human brain malignant glioma tissue, Human bladder carcinoma tissue;
Primary antibody: 217272(CDK5RAP3 Antibody) at dilution 1/449;
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
Exposure time: 40 seconds