



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **CFAP298 RABBIT PAB**


货号: S218409 产品全名: CFAP298 兔多抗 基因符号 Kur; FBB18; CILD26; C21orf48; C21orf59 UNIPROT ID: P57076 (Gene Accession - BC000709) 背景: This gene encodes a protein that plays a critical role in dynein arm assembly and motile cilia function. Mutations in this gene result in primary ciliary dyskinesia. Naturally occuring readthrough transcription occurs from this locus to the downstream t-complex 10 like (TCP10L) gene. [provided by RefSeq, Apr 2017] 抗原: Fusion protein of human CFAP298 经过测试的应用: ELISA, WB, IHC 推荐稀释比: IHC: 25-100;WB: 200-1000;ELISA: 5000-10000 种属反应性: Rabbit 克隆性: Rabbit Polyclonal 亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat 成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol 研究领域: Signal Transduction 储存和运输: Store at -20°C. Avoid repeated freezing and thawing





Immunohistochemistry analysis of paraffin embedded Human prostate cancer tissue using 218409(CFAP298 Antibody) at a dilution of 1/30(Cytoplasm).

In comparision with the IHC on the left, the same paraffin-embedded Human prostate cancer tissue is first treated with the fusion protein and then with 218409(Anti-CFAP298 Antibody) at dilution 1/30.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: Human testis tissue lysate; Primary antibody: 218409(CFAP298 Antibody) at dilution 1/250; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 1 second