

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

CFLAR RABBIT PAB

货号: S220553

产品全名: CFLAR 兔多抗

基因符号 CASH; FLIP; MRIT; CLARP; FLAME; Casper; FLAMEI; c-FLIP; FLAME-1; I-FLICE; c-FLIPL; c-FLIPR; c-FLIPS; CASP8AP1

UNIPROT ID: 015519 (Gene Accession - NP_003870)

背景: The protein encoded by this gene is a regulator of apoptosis and is structurally similar to caspase-8. However, the encoded protein lacks caspase activity and appears to be itself cleaved into two peptides by caspase-8. Several transcript variants encoding different isoforms have been found for this gene, and partial evidence for several more variants exists.

抗原: Synthetic peptide of human CFLAR

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 200-1000;ELISA: 1000-2000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction, Cancer

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human ovarian cancer tissue using 220553(CFLAR Antibody) at a dilution of 1/20(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human ovarian cancer tissue is first treated with the synthetic peptide and then with 220553(Anti-CFLAR Antibody) at dilution 1/20.



The image on the left is immunohistochemistry of paraffinembedded Human colon cancer tissue using 220553(Anti-CFLAR Antibody) at a dilution of 1/20.



In comparision with the IHC on the left, the same paraffin-embedded Human colon cancer tissue is first treated with synthetic peptide and then with D261685(Anti-CFLAR Antibody) at dilution 1/20.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: Mouse heart tissue; Primary antibody: 220553(CFLAR Antibody) at dilution 1/200; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 4 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010