

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **CLIC1 RABBIT PAB**

货号: S217629

产品全名: CLIC1 兔多抗 基因符号 G6; NCC27

UNIPROT ID: 000299 (Gene Accession - BC064527)

背景: Chloride channels are a diverse group of proteins that regulate fundamental cellular processes including stabilization of cell membrane potential, transepithelial transport, maintenance of intracellular pH, and regulation of cell volume. Chloride intracellular channel 1 is a member of the p64 family; the protein localizes principally to the cell nucleus and exhibits both nuclear and plasma membrane chloride ion channel activity.

抗原: Fusion protein of human CLIC1 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-200;WB: 200-1000;ELISA: 1000-2000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Metabolism, Signal Transduction

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 217629(CLIC1 Antibody) at a dilution of 1/30(Cytoplasm).




In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the fusion protein and then with 217629(Anti-CLIC1 Antibody) at dilution 1/30.



The image on the left is immunohistochemistry of paraffinembedded Human breast cancer tissue using 217629(Anti-CLIC1 Antibody) at a dilution of 1/30.



In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with fusion protein and then with D222732(Anti-CLIC1 Antibody) at dilution 1/30.



Gel: 10%SDS-PAGE, Lysate: 40 µg; Lane: Human testis tissue; Primary antibody: 217629(CLIC1 Antibody) at dilution 1/400; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 1 minute



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010