

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

COX7B RABBIT PAB

货号: S216277

产品全名: COX7B 兔多抗

基因符号 APLCC

UNIPROT ID: P24311 (Gene Accession - BC018386)

背景: Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes subunit VIIb, which is highly similar to bovine COX VIIb protein and is found in all tissues. This gene may have several pseudogenes on chromosomes 1, 2, 20 and 22.

抗原: Fusion protein of human COX7B

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 100-300;WB: 500-2000;ELISA: 2000-5000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification

种属反应性: Human, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Metabolism

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




cancer tissue using 216277(COX7B Antibody) at a dilution of 1/70(Cytoplasm).



Immunohistochemistry analysis of paraffin embedded Human colon In comparision with the IHC on the left, the same paraffin-embedded Human colon cancer tissue is first treated with the fusion protein and then with 216277(Anti-COX7B Antibody) at dilution 1/70.



The image on the left is immunohistochemistry of paraffinembedded Human gasrtic cancer tissue using 216277(Anti-COX7B Antibody) at a dilution of 1/70.



In comparision with the IHC on the left, the same paraffin-embedded Human gasrtic cancer tissue is first treated with fusion protein and then with D220115(Anti-COX7B Antibody) at dilution 1/70.

40-25-4.6

kDa

Gel: 10+15%SDS-PAGE, Lysate: 50 µg; Lane: 293T cells; Primary antibody: 216277(COX7B Antibody) at dilution 1/800; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 40 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010