

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

CSF2RA RABBIT PAB

产品全名: CSF2RA 兔多抗 基因符号 GMR; CDI16; CSF2R; SMDP4; CDw116; CSF2RX; CSF2RY; GMCSFR; CSF2RAX; CSF2RAY; GM-CSF-R-alpha UNIPROT ID: P15509 (Gene Accession - NP_006131) 背景: The protein encoded by this gene is the alpha subunit of the heterodimeric receptor for colony stimulating factor 2, a cytokine which controls the production, differentiation, and function of granulocytes and macrophages. The encoded protein is a member of the cytokine family of receptors. This gene is found in the pseudoautosomal region (PAR) of the X and Y chromosomes. Multiple transcript variants encoding different isoforms have been found for this gene, with some of the isoforms being membrane-bound and others being soluble. 抗原: Synthetic peptide of human CSF2RA 经过测试的应用: ELISA, WB, IHC 推荐稀释比: IHC: 50-200;WB: 500-2000;ELISA: 2000-5000 种属反应性: Rabbit 克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG

纯化: Antigen affinity purification

种属反应性: Human

货号: S219827

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Immunology, Stem Cells

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human cervical cancer tissue using 219827(CSF2RA Antibody) at a dilution of Human cervical cancer tissue is first treated with the synthetic 1/35(Cytoplasm).




The image on the left is immunohistochemistry of paraffinembedded Human thyroid cancer tissue using 219827(Anti-CSF2RA Antibody) at a dilution of 1/35.



In comparision with the IHC on the left, the same paraffin-embedded



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with synthetic peptide and then with D260442(Anti-CSF2RA Antibody) at dilution 1/35.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-2: Human placenta tissue, HepG2 cells; Primary antibody: 219827(CSF2RA Antibody) at dilution 1/250; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 3 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010