

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## CYP2E1 RABBIT PAB

货号: S217320

产品全名: CYP2EI 兔多抗

基因符号 CPE1; CYP2E; P450-J; P450C2E

UNIPROT ID: P05181 (Gene Accession - BC067435)

背景: This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and is induced by ethanol, the diabetic state, and starvation. The enzyme metabolizes both endogenous substrates, such as ethanol, acetone, and acetal, as well as exogenous substrates including benzene, carbon tetrachloride, ethylene glycol, and nitrosamines which are premutagens found in cigarette smoke. Due to its many substrates, this enzyme may be involved in such varied processes as gluconeogenesis, hepatic cirrhosis, diabetes, and cancer.

抗原: Fusion protein of human CYP2E1

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-200;WB: 1000-5000;ELISA: 5000-10000

种属反应性: Rabbit


克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat


成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Metabolism, Cardiovascular


储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 217320(CYP2E1 Antibody) at a dilution of 1/80(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the fusion protein and then with 217320(Anti-CYP2EI Antibody) at dilution 1/80.



Gel: 8%SDS-PAGE, Lysate:40 µg;

Lane 1-3: Mouse fetal liver tissue, Mouse kidney tissue, Mouse liver

tissue lysates:

Primary antibody: 217320(CYP2E1 Antibody) at dilution 1/1150;

Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000

dilution;

Exposure time: 1 second



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010