

CYSTATIN C (3C8) MOUSE MAB

货号: N261311

产品全名: Cystatin C (3C8) 小鼠单抗

基因符号 CST3; Cystatin-C; Cystatin-3; Gamma-trace; Neuroendocrine basic polypeptide; Post-gamma-globulin

UNIPROT ID: P01034

背景: Cystatin C is a 14 kDa member of the Cystatin superfamily of cysteine protease inhibitors. Most cell types secrete Cystatin C. Cystatin C inhibits cathepsins, and thereby may function as a tumor suppressor by inhibiting cathepsin mediated tumor cell invasion. In addition, this tumor suppressor function can also be attributed to Cystatin C's ability to antagonize TGF- β 1 signaling.

抗原: Purified recombinant protein expressed in E.coli.

经过测试的应用: WB,ELISA

推荐稀释比: WB: 1/500-1/1000 ELISA: 1/10000

种属反应性: Mouse

克隆性: Mouse Monoclonal

克隆编号: 3C8-2E9-8C5

分子量: Calculated MW: 16 kDa; Observed MW: 16 kDa

亚型: IgG1

纯化: Affinity Purified

种属反应性: Human

成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

研究领域: Tags & Cell Markers

储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Western blot analysis of Cystatin C (3C8) in HeLa, A549, HL-60, U2OS, C6 lysates using Cystatin C (3C8) antibody.



Product Description

Pioneering GTPase and Oncogene Product Development since 2010
