

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **DDX56 RABBIT PAB**

货号: S221885

产品全名: DDX56 兔多抗 基因符号 DDX21; DDX26; NOH61

UNIPROT ID: Q9NY93 (Gene Accession - NP\_061955)

背景: This gene encodes a member of the DEAD box protein family. DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. The protein encoded by this gene shows ATPase activity in the presence of polynucleotides and associates with nucleoplasmic 65S preribosomal particles. This gene may be involved in ribosome synthesis, most likely during assembly of the large 60S ribosomal subunit. Multiple transcript variants encoding different isoforms have been found for this gene.

抗原: Synthetic peptide of human DDX56

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 20-100;WB: 200-1000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal


亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification

种属反应性: Human


成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Epigenetics and Nuclear Signaling


储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human lung cancer tissue using 221885(DDX56 Antibody) at a dilution of 1/25(Nucleus).



In comparision with the IHC on the left, the same paraffin-embedded Human lung cancer tissue is first treated with the synthetic peptide and then with 221885(Anti-DDX56 Antibody) at dilution 1/25.



Gel: 8%SDS-PAGE, Lysate: 40 µg;

Lane 1-3: HEPG2, 293T and Hela cell lysates;

Primary antibody: 221885(DDX56 Antibody) at dilution 1/400; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;

Exposure time: 7 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010