

EDN3 RABBIT PAB

货号: S216503

产品全名: EDN3 兔多抗

基因符号 ET3; ET-3; WS4B; HSCR4; PPET3

UNIPROT ID: P14138 (Gene Accession - BC008876)

背景: The protein encoded by this gene is a member of the endothelin family. Endothelins are endothelium-derived vasoactive peptides involved in a variety of biological functions. The active form of this protein is a 21 amino acid peptide processed from the precursor protein. The active peptide is a ligand for endothelin receptor type B (EDNRB). The interaction of this endothelin with EDNRB is essential for development of neural crest-derived cell lineages, such as melanocytes and enteric neurons. Mutations in this gene and EDNRB have been associated with Hirschsprung disease (HSCR) and Waardenburg syndrome (WS), which are congenital disorders involving neural crest-derived cells. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript variants encoding different isoforms.

抗原: Fusion protein of human EDN3

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-200;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human, Mouse


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Cardiovascular, Neuroscience


储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 216503(EDN3 Antibody) at a dilution of 1/60 (Secreted).



In comparison with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the fusion protein and then with 216503(Anti-EDN3 Antibody) at dilution 1/60.




The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using 216503(Anti-EDN3 Antibody) at a dilution of 1/60.



In comparison with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with fusion protein and then with D220612(Anti-EDN3 Antibody) at dilution 1/60.

kDa



Gel: 12%SDS-PAGE, Lysate: 40 µg;
 Lane: Mouse eye tissue lysate;
 Primary antibody: 216503(EDN3 Antibody) at dilution 1/800;
 Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;
 Exposure time: 30 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010
