

EIF3K RABBIT PAB

货号: S219018

产品全名: EIF3K 兔多抗

基因符号 M9; ARG134; PLAC24; PTD001; EIF3S12; HSPC029; MSTP001; PLAC-24; PRO1474; EIF3-p28

UNIPROT ID: Q9UBQ5 (Gene Accession - BC001031)

背景: The 700-kD eukaryotic translation initiation factor-3 (eIF3) is the largest eIF and contains at least 12 subunits, including EIF2S12. eIF3 plays an essential role in translation by binding directly to the 40S ribosomal subunit and promoting formation of the 40S preinitiation complex.

抗原: Fusion protein of human EIF3K

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-300; WB: 500-2000; ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human, Mouse


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Epigenetics and Nuclear Signaling, Neuroscience

储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 219018(EIF3K Antibody) at a dilution of 1/55(Cytoplasm).



In comparison with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the fusion protein and then with 219018(Anti-EIF3K Antibody) at dilution 1/55.



Gel: 12%SDS-PAGE, Lysate: 40 µg;

Lane: HT29 cell lysate;

Primary antibody: 219018(EIF3K Antibody) at dilution 1/400;

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;

Exposure time: 10 seconds