

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **EPSTI1 RABBIT PAB**

货号: S217406 产品全名: EPSTII 兔多抗

基因符号 BRESII

UNIPROT ID: Q96J88 (Gene Accession - BC023660)

背景: The protein encoded by this gene has been shown to promote tumor invasion and metastasis in some invasive cancer cells when overexpressed. Expression of this gene has been shown to be upregulated by direct binding of the Kruppel like factor 8 protein to promoter sequences. The translated protein interacts with the amino terminal region of the valosin containing protein gene product, resulting in the nuclear translocation of the nuclear factor kappa B subunit 1 gene product, and activation of target genes. Overexpression of this gene has been observed in some breast cancers and in some individuals with systemic lupus erythematosus (SLE). [provided by RefSeq, Sep 2016]

抗原: Fusion protein of human EPSTII 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 100-200;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit 克隆性: Rabbit Polyclonal


亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification

种属反应性: Human


成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Cancer


储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human breast cancer tissue using 217406(EPSTII Antibody) at a dilution of 1/160(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with the fusion protein and then with 217406(Anti-EPSTII Antibody) at dilution 1/160.



Gel: 8%SDS-PAGE, Lysate: 40 μg; Lane: HepG2 cell lysate;

Primary antibody: 217406(EPSTII Antibody) at dilution 1/1000;

Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000

dilution;

Exposure time: 1 minute