

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

FCRLA RABBIT PAB

货号: S219354

产品全名: FCRLA 兔多抗

基因符号 FCRL; FCRX; FREB; FCRL1; FCRLX; FCRLb; FCRLd; FCRLe; FCRLM1; FCRLc1; FCRLc2

UNIPROT ID: Q7L513 (Gene Accession - BC006521)

背景: This gene encodes a protein similar to receptors for the Fc fragment of gamma immunoglobulin (IgG). These receptors, referred to as FCGRs, mediate the destruction of IgG-coated antigens and of cells induced by antibodies. This encoded protein is selectively expressed in B cells, and may be involved in their development. This protein may also be involved in the development of lymphomas. Multiple alternatively spliced transcript variants that encode different protein isoforms have been described for this gene.

抗原: Fusion protein of human FCRLA

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 100-200;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification

种属反应性: Human

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Immunology

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description

Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human colorectal cancer tissue using 219354(FCRLA Antibody) at a dilution of 1/90(Cytoplasm).


In comparision with the IHC on the left, the same paraffin-embedded Human colorectal cancer tissue is first treated with the fusion protein and then with 219354(Anti-FCRLA Antibody) at dilution 1/90.



The image on the left is immunohistochemistry of paraffinembedded Human prostate cancer tissue using 219354(Anti-FCRLA Antibody) at a dilution of 1/90.



In comparision with the IHC on the left, the same paraffin-embedded Human prostate cancer tissue is first treated with fusion protein and then with D226511(Anti-FCRLA Antibody) at dilution 1/90.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: Ramos cell lysate; Primary antibody: 219354(FCRLA Antibody) at dilution 1/750; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;

Exposure time: 1 minute



Product Description

Pioneering GTPase and Oncogene Product Development since 2010