

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

FKBP1A RABBIT PAB

货号: S218942 产品全名: FKBP1A 兔多抗 基因符号 FKBP1; PKC12; PKC12; FKBP12; PPIASE; FKBP-12; FKBP-1A UNIPROT ID: P62942 (Gene Accession - BC005147)

背景: The protein encoded by this gene is a member of the immunophilin protein family, which play a role in immunoregulation and basic cellular processes involving protein folding and trafficking. The protein is a cis-trans prolyl isomerase that binds the immunosuppressants FK506 and rapamycin. It interacts with several intracellular signal transduction proteins including type I TGF-beta receptor. It also interacts with multiple intracellular calcium release channels, and coordinates multi-protein complex formation of the tetrameric skeletal muscle ryanodine receptor. In mouse, deletion of this homologous gene causes congenital heart disorder known as noncompaction of left ventricular myocardium. Multiple alternatively spliced variants, encoding the same protein, have been identified. The human genome contains five pseudogenes related to this gene, at least one of which is transcribed.

抗原: Fusion protein of human FKBP1A

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 40-200;WB: 200-1000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG

纯化: Antigen affinity purification


种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Immunology


储存和运输: Store at -20°C. Avoid repeated freezing and thawing





tissue using 218942(FKBP1A Antibody) at a dilution of

Immunohistochemistry analysis of paraffin embedded Human tonsil In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the fusion protein and then with 218942(Anti-FKBP1A Antibody) at dilution 1/55



Gel: 12%SDS-PAGE, Lysate: 40 µg;

Lane 1-3: Mouse brain tissue, Human fetal brain tissue and Jurkat cell lvsates:

Primary antibody: 218942(FKBP1A Antibody) at dilution 1/250; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 2 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010