

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

FTO RABBIT PAB

货号: \$217458 产品全名: FTO 兔多抗

基因符号 GDFD; ALKBH9; BMIQ14

UNIPROT ID: Q9C0B1 (Gene Accession - BC003583)

背景: This gene is a nuclear protein of the AlkB related non-haem iron and 2-oxoglutarate-dependent oxygenase superfamily but the exact physiological function of this gene is not known. Other non-heme iron enzymes function to reverse alkylated DNA and RNA damage by oxidative demethylation. Studies in mice and humans indicate a role in nervous and cardiovascular systems and a strong association with body mass inde,x obesity risk, and type 2 diabetes. [provided by RefSeq, Jul 2011]

抗原: Fusion protein of human FTO 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-200;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification

种属反应性: Human

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Metabolism, Neuroscience, Cardiovascular

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human prostate cancer tissue using 217458(FTO Antibody) at a dilution of 1/80(Cytoplasm and Nucleus).



In comparision with the IHC on the left, the same paraffin-embedded Human prostate cancer tissue is first treated with the fusion protein and then with 217458(Anti-FTO Antibody) at dilution 1/80.



The image on the left is immunohistochemistry of paraffinembedded Human tonsil tissue using 217458(Anti-FTO Antibody) at a Human tonsil tissue is first treated with fusion protein and then with dilution of 1/80.



In comparision with the IHC on the left, the same paraffin-embedded D222388(Anti-FTO Antibody) at dilution 1/80.

kDa 130-100-70 **—** 55 **—** 35 **—** 25 — 15 **—**

Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: HepG2 cell lysate; Primary antibody: 217458(FTO Antibody) at dilution 1/1150; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution; Exposure time: 90 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010