

GLYR1 RABBIT PAB

货号: S214364

产品全名: GLYR1 兔多抗

基因符号 NP60; BMO45; HIBDL; N-PAC

UNIPROT ID: Q49A26 (Gene Accession - NP_115958)

背景: Nucleosome-destabilizing factor that is recruited to genes during transcriptional activation (PubMed:29759984). Facilitates Pol II transcription through nucleosomes (PubMed:29759984). Binds DNA (in vitro) (PubMed:29759984). Recognizes and binds trimethylated 'Lys-36' of histone H3 (H3K36me3) (PubMed:20850016). Promotes KDM1B demethylase activity (PubMed:23260659). Stimulates the acetylation of 'Lys-56' of nucleosomal histone H3 (H3K56ac) by EP300 (PubMed:29759984). Regulates p38 MAP kinase activity by mediating stress activation of p38alpha/MAPK14 and specifically regulating MAPK14 signaling (PubMed:16352664). Indirectly promotes phosphorylation of MAPK14 and activation of ATF2 (PubMed:16352664). The phosphorylation of MAPK14 requires upstream activity of MAP2K4 and MAP2K6 (PubMed:16352664). Putative oxidoreductase (PubMed:23260659).

抗原: Synthetic peptide of human GLYR1

经过测试的应用: ELISA, IHC

推荐稀释比: IHC: 50-200; ELISA: 2000-5000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human, Mouse, Rat


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Signal Transduction, Epigenetics and Nuclear Signaling


储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human cervical cancer tissue using 214364 (GLYR1 Antibody) at a dilution of 1/30 (Nucleus).



In comparison with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with the synthetic peptide and then with 214364 (Anti-GLYR1 Antibody) at dilution 1/30.



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using 214364 (Anti-GLYR1 Antibody) at a dilution of 1/30.



In comparison with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with synthetic peptide and then with D161717 (Anti-GLYR1 Antibody) at dilution 1/30.