

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **GPD2 RABBIT PAB**

货号: S220584

产品全名: GPD2 兔多抗

基因符号 GDH2; GPDM; mGPDH

UNIPROT ID: P43304 (Gene Accession - NP\_000399)

背景: The protein encoded by this gene localizes to the inner mitochondrial membrane and catalyzes the conversion of glycerol-3-phosphate to dihydroxyacetone phosphate, using FAD as a cofactor. Along with GDPI, the encoded protein constitutes the glycerol phosphate shuttle, which reoxidizes NADH formed during glycolysis. Two transcript variants encoding the same protein have been found for this gene.

抗原: Synthetic peptide of human GPD2

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 20-100;WB: 500-2000;ELISA: 2000-5000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Metabolism, Cancer

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human esophagus cancer tissue using 220584(GPD2 Antibody) at a dilution of 1/20(Cytoplasm).




In comparision with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with the synthetic peptide and then with 220584(Anti-GPD2 Antibody) at dilution 1/20.



The image on the left is immunohistochemistry of paraffinembedded Human thyroid cancer tissue using 220584(Anti-GPD2 Antibody) at a dilution of 1/20.



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with synthetic peptide and then with D261724(Anti-GPD2 Antibody) at dilution 1/20.



Gel: 6%SDS-PAGE, Lysate: 40 µg; Lane 1-3: A172 cells, 231 cells, Hela cells; Primary antibody: 220584(GPD2 Antibody) at dilution 1/200; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 30 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010