

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

HCK RABBIT PAB

货号: S210354 产品全名: HCK 兔多抗

基因符号 JTK9; p59Hck; p61Hck

UNIPROT ID: P08631 (Gene Accession - BC014435)

背景: The protein encoded by this gene is a member of the Src family of tyrosine kinases. This protein is primarily hemopoietic, particularly in cells of the myeloid and B-lymphoid lineages. It may help couple the Fc receptor to the activation of the respiratory burst. In addition, it may play a role in neutrophil migration and in the degranulation of neutrophils. Multiple isoforms with different subcellular distributions are produced due to both alternative splicing and the use of alternative translation initiation codons, including a non-AUG (CUG) codon.

抗原: Fusion protein of human HCK 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 200-1000;ELISA: 2000-5000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction, Immunology

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 210354(HCK Antibody) at a dilution of 1/20(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the fusion protein and then with 210354(Anti-HCK Antibody) at dilution 1/20.



The image on the left is immunohistochemistry of paraffinembedded Human liver cancer tissue using 210354(Anti-HCK Antibody) at a dilution of 1/20.



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with fusion protein and then with D120736(Anti-HCK Antibody) at dilution 1/20.

%%%%Gel: 8%SDS-PAGE, Lysate: 40 μg;
Lane: A549 cells;

Gel: 8%SDS-PAGE, Lysate: 40 μg; Lane: A549 cells; Primary antibody: 210354(HCK Antibody) at dilution 1/450; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 15 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010