

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## HSD3B7 RABBIT PAB

货号: S217521 产品全名: HSD3B7 兔多抗 基因符号 CBASI; PFIC4; SDR1IE3 UNIPROT ID: Q9H2F3 (Gene Accession - BC004929)

背景: This gene encodes an enzyme which is involved in the initial stages of the synthesis of bile acids from cholesterol and a member of the short-chain dehydrogenase/reductase superfamily. The encoded protein is a membrane-associated endoplasmic reticulum protein which is active against 7-alpha hydrosylated sterol substrates. Mutations in this gene are associated with a congenital bile acid synthesis defect which leads to neonatal cholestasis, a form of progressive liver disease. Multiple transcript variants encoding different isoforms have been found for this aene.

抗原: Fusion protein of human HSD3B7

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-200;WB: 500-2000;ELISA: 2000-5000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG

纯化: Antigen affinity purification

种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 217521(HSD3B7 Antibody) at a dilution of 1/40(Cytoplasm).



The image on the left is immunohistochemistry of paraffinembedded Human liver cancer tissue using 217521(Anti-HSD3B7 Antibody) at a dilution of 1/40.



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the fusion protein and then with 217521(Anti-HSD3B7 Antibody) at dilution 1/40.



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with fusion protein and then with D222512(Anti-HSD3B7 Antibody) at dilution 1/40.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: Skov3 cells; Primary antibody: 217521(HSD3B7 Antibody) at dilution 1/400; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 10 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010