

ICOSLG RABBIT PAB

货号: S213390

产品全名: ICOSLG 兔多抗

基因符号 B7h; B7H2; GL50; B7-H2; B7RP1; CD275; ICOSL; LICOS; B7RP-1; ICOS-L

UNIPROT ID: O75144 (Gene Accession - NP_056074)

背景: Enables identical protein binding activity. Predicted to be involved in T cell receptor signaling pathway and positive regulation of interleukin-4 production. Located in cytoplasmic ribonucleoprotein granule and plasma membrane.

抗原: Synthetic peptide of human ICOSLG

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-200;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Immunology

储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human esophagus cancer tissue using 213390(ICOSLG Antibody) at a dilution of 1/50(Cytoplasm and Cell membrane).



Gel: 12%SDS-PAGE, Lysate: 40 µg;

Lane 1-2: A172 and HUVEC cell lysates;

Primary antibody: 213390(ICOSLG Antibody) at dilution 1/650;

Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;

Exposure time: 2 minutes

✖ In comparison with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with the synthetic peptide and then with 213390(Anti-ICOSLG Antibody) at dilution 1/50.