Product Description

Pioneering GTPase and Oncogene Product Development since 2010

IL17RC RABBIT PAB

货号: S216301
产品全名: IL17RC 兔多抗
基因符号 CANDF9; IL17RL; IL17-RL
UNIPROT ID: Q8NAC3 (Gene Accession – BC006411)
背景: This gene encodes a single-pass type I membrane protein that shares similarity with the interleukin-17 receptor (IL-17RA). Unlike IL-17RA, which is predominantly expressed in hemopoietic cells, and binds with high affinity to only IL-17A, this protein is expressed in nonhemopoietic tissues, and binds both IL-17A and IL-17F with similar affinities. The proinflammatory cytokines, IL-17A and IL-17F, have been implicated in the progression of inflammatory and autoimmune diseases. Multiple alternatively spliced transcript variants encoding different isoforms have been detected for this gene, and it has been proposed that soluble, secreted proteins lacking transmembrane and intracellular domains may function as extracellular antagonists to cytokine signaling.
抗原: Fusion protein of human IL17RC
经过测试的应用: ELISA, IHC
推荐稀释比: IHC: 50-200; ELISA: 5000-10000
种属反应性: Rabbit
克隆性： Rabbit Polyclonal
亚型: Immunogen-specific rabbit IgG
纯化: Antigen affinity purification
种属反应性: Human
成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol
研究领域: Immunology
储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human colorectal cancer tissue using 216301(IL17RC Antibody) at a dilution of 1/55(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human colorectal cancer tissue is first treated with the fusion protein and then with 216301(Anti-IL17RC Antibody) at dilution 1/55.



The image on the left is immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using 216301(Anti-IL17RC Antibody) at a dilution of 1/55.



In comparision with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with fusion protein and then with D220168(Anti-IL17RC Antibody) at dilution 1/55.