L1RE1 RABBIT PAB

货号: S221817
产品全名: L1RE1 兔多抗
基因符号 L1.2; LRE1
UNIPROT ID: Q9UN81
背景: Nucleic acid-binding protein which is essential for retrotransposition of LINE-1 elements in the genome. May function as a nucleic acid chaperone binding its own transcript and therefore preferentially mobilizing the transcript from which they are encoded.
抗原: Synthetic peptide of human L1RE1
经过测试的应用: ELISA, WB, IHC
推荐稀释比: IHC: 25-100;WB: 200-1000;ELISA: 5000-10000
种属反应性: Rabbit
克隆性： Rabbit Polyclonal
亚型: Immunogen-specific rabbit IgG
纯化: Antigen affinity purification
种属反应性: Human
成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol
研究领域: Epigenetics and Nuclear Signaling
储存和运输: Store at -20°C. Avoid repeated freezing and thawing

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC APPLICATIONS

Product Description

Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 221817(L1RE1 Antibody) at a dilution of 1/20(Cytoplasm or Nucleus).



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the synthetic peptide and then with 221817(Anti-L1RE1 Antibody) at dilution 1/20.



The image on the left is immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using 221817(Anti-L1RE1 Antibody) at a dilution of 1/20.



In comparision with the IHC on the left, the same paraffin-embedded Human ovarian cancer tissue is first treated with synthetic peptide and then with D263576(Anti-L1RE1 Antibody) at dilution 1/20.



Gel: 8%SDS-PAGE, Lysate: 40 µg;
Lane 1-2: A431 and 293T cell lysates;
Primary antibody: 221817(L1RE1 Antibody) at dilution 1/200;
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
Exposure time: 10 seconds

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC APPLICATIONS

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC APPLICATIONS