

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **LAMININ GAMMA 2 RABBIT PAB**

货号: N225036

产品全名: Laminin gamma 2 兔多抗

基因符号 LAMC2; LAMB2T; LAMNB2; Laminin subunit gamma-2; Cell-scattering factor 140 kDa subunit; CSF 140 kDa subunit; Epiligrin subunit gamma; Kalinin subunit gamma; Kalinin/nicein/epiligrin 100 kDa subunitLadsin 140 kDa subunit; Laminin B2t chain; Laminin-5 subunit gamma; Large adhesive scatter factor 140 kDa subunit; Nicein subunit gamma

**UNIPROT ID:** Q13753

背景: Binding to cells via a high affinity receptor, laminin is thought to mediate the attachment, migration and organization of cells into tissues during embryonic development by interacting with other extracellular matrix components.

抗原: The antiserum was produced against synthesized peptide derived from the C-terminal region of human LAMC2. AA range:1021-1070

经过测试的应用: WB,IHC-P,ELISA

推荐稀释比: WB: 1/500-1/1000 IHC: 1/50-1/100 ELISA: 1/10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

分子量: Calculated MW: 131 kDa; Observed MW: 135 kDa

亚型: IgG

纯化: Affinity Purified 种属反应性: Human,Rat

成分: PBS (without Mg2+ and Cg2+), pH 7.3 containing 50% glycerol,

0.5% BSA and 0.02% sodium azide


研究领域: Cardiovascular

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin-embedded Human liver using Laminin gamma 2 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of Laminin gamma 2 in K562 lysates using Laminin gamma 2 antibody.



paraffin-embedded Human lung using Laminin gamma 2 antibody. using Laminin gamma 2 High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemistry analysis of Immunohistochemistry analysis of paraffin-embedded Human lung antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.