

LRIG1 RABBIT PAB

货号: S219894

产品全名: LRIG1 兔多抗

基因符号: LIG1; LIG-1

UNIPROT ID: Q96JA1 (Gene Accession - NP_056356)

背景: Predicted to act upstream of or within several processes, including innervation; otolith morphogenesis; and sensory perception of sound. Predicted to be located in plasma membrane. Predicted to be active in extracellular matrix and extracellular space.

抗原: Synthetic peptide of human LRIG1

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 100-300;WB: 1000-5000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human, Mouse


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Cancer


储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human cervical cancer tissue using 219894(LRIG1 Antibody) at a dilution of 1/95(Cytoplasm).



In comparison with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with the synthetic peptide and then with 219894(Anti-LRIG1 Antibody) at dilution 1/95.




The image on the left is immunohistochemistry of paraffin-embedded Human prostate cancer tissue using 219894(Anti-LRIG1 Antibody) at a dilution of 1/95.



In comparison with the IHC on the left, the same paraffin-embedded Human prostate cancer tissue is first treated with synthetic peptide and then with D260563(Anti-LRIG1 Antibody) at dilution 1/95.

kDa



Gel: 6%SDS-PAGE, Lysate: 40 µg;
Lane: Mouse skeletal muscle tissue lysate;
Primary antibody: 219894(LRIG1 Antibody) at dilution 1/1300;
Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;
Exposure time: 10 seconds

