

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

MAP1LC3B RABBIT PAB

货号: S221803

产品全名: MAPILC3B 兔多抗

基因符号 LC3B; ATG8F; MAPILC3B-a; MAPIA/IBLC3
UNIPROT ID: Q9GZQ8 (Gene Accession - NP_073729)

背景: The product of this gene is a subunit of neuronal microtubule-associated MAPIA and MAPIB proteins, which are involved in microtubule assembly and important for neurogenesis. Studies on the rat homolog implicate a role for this gene in autophagy, a process that involves the bulk degradation of cytoplasmic component.

抗原: Synthetic peptide of human MAPILC3B

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 20-100;WB: 200-1000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat


成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Signal Transduction, Cancer, Neuroscience, Cardiovascular 储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 221803(MAPILC3B Antibody) at a dilution of 1/25(Cytoplasm or Nucleus).




In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the synthetic peptide and then with 221803(Anti-MAPILC3B Antibody) at dilution 1/25.



The image on the left is immunohistochemistry of paraffinembedded Human prostate cancer tissue using 221803(Anti-MAPILC3B Antibody) at a dilution of 1/25.



In comparision with the IHC on the left, the same paraffin-embedded Human prostate cancer tissue is first treated with synthetic peptide and then with D263557(Anti-MAPILC3B Antibody) at dilution 1/25.



Gel: 12%SDS-PAGE, Lysate: 40 µg; Lane 1-2: HEPG2 cell and Human fetal brain tissue lysates; Primary antibody: 221803(MAP1LC3B Antibody) at dilution 1/300; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 5 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010