

MOCOS RABBIT PAB

货号: S219537

产品全名: MOCOS 兔多抗

基因符号: MCS; MOS; HMCS

UNIPROT ID: Q96EN8 (Gene Accession - BC012079)

背景: This gene encodes an enzyme that sulfurates the molybdenum cofactor which is required for activation of the xanthine dehydrogenase (XDH) and aldehyde oxidase (AO) enzymes. XDH catalyzes the conversion of hypoxanthine to uric acid via xanthine, as well as the conversion of allopurinol to oxypurinol, and pyrazinamide to 5-hydroxy pyrazinamide. Mutations in this gene cause the metabolic disorder classical xanthinuria type II which is characterized by the loss of XDH/XO and AO enzyme activity, decreased levels of uric acid in the urine, increased levels of xanthine and hypoxanthine in the serum and urine, formation of xanthine stones in the urinary tract, and myositis due to tissue deposition of xanthine.

抗原: Fusion protein of human MOCOS

经过测试的应用: ELISA, IHC

推荐稀释比: IHC: 50-100; ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human

成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Cell Biology

储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 219537(MOCOS Antibody) at a dilution of 1/50(Cytoplasm).

In comparison with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the fusion protein and then with 219537(Anti-MOCOS Antibody) at dilution 1/50.