

NAA10 RABBIT PAB

货号: S219030

产品全名: NAA10 兔多抗

基因符号 TE2; ARD1; NATD; ARD1A; ARD1P; OGDNS; hARD1; DXS707; MCOPSI

UNIPROT ID: P41227 (Gene Accession - BC000308)

背景: N-alpha-acetylation is among the most common post-translational protein modifications in eukaryotic cells. This process involves the transfer of an acetyl group from acetyl-coenzyme A to the alpha-amino group on a nascent polypeptide and is essential for normal cell function. This gene encodes an N-terminal acetyltransferase that functions as the catalytic subunit of the major amino-terminal acetyltransferase A complex. Mutations in this gene are the cause of Ogden syndrome. Alternate splicing results in multiple transcript variants.

抗原: Fusion protein of human NAA10

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-300;WB: 200-1000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human, Mouse


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Metabolism, Epigenetics and Nuclear Signaling


储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 219030(NAA10 Antibody) at a dilution of 1/55(Cytoplasm or Nucleus).




In comparison with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the fusion protein and then with 219030(Anti-NAA10 Antibody) at dilution 1/55.



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using 219030(Anti-NAA10 Antibody) at a dilution of 1/55.



In comparison with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with fusion protein and then with D225680(Anti-NAA10 Antibody) at dilution 1/55.



Gel: 12%SDS-PAGE, Lysate: 40 µg;
Lane 1-2: HeLa and 293T cell lysates;
Primary antibody: 219030(NAA10 Antibody) at dilution 1/400;
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
Exposure time: 10 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010
