

## NF-KB P65 RABBIT PAB

货号: N225543

产品全名: NF-KB p65 兔多抗

基因符号 NFKB3; RELA; TF65; Transcription factor p65; p65; NFkB

**UNIPROT ID:** Q04206

背景: NFKB1 (MIM 164011) or NFKB2 (MIM 164012) is bound to REL (MIM 164910), RELA, or RELB (MIM 604758) to form the NFKB complex. The p50 (NFKB1)/p65 (RELA) heterodimer is the most abundant form of NFKB. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA, MIM 164008 or NFKBIB, MIM 604495), which inactivate NFKB by trapping it in the cytoplasm.

抗原: A synthesized peptide derived from human NF-kB p65

经过测试的应用: WB,IHC-P,ICC/IF,IP,FC

推荐稀释比: WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/20 FC: 1/50-1/100

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

分子量: Calculated MW: 60 kDa; Observed MW: 65 kDa

亚型: IgG


纯化: Affinity Chromatography

种属反应性: Human,Mouse


成分: PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

研究领域: Cell Biology


储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Western blot analysis of NF-κB p65 expression in HeLa lysates using NF-κB p65R antibody



Immunofluorescence analysis of NF-κB p65 in HT-1080 using NF-κB p65 antibody.



Western blot analysis of NF-κB p65 in MCF-7, HeLa, Molt4, Raji and Jurkat lysates using NF-κB p65 antibody.



Immunohistochemistry analysis of paraffin-embedded Human transitional carcinoma of bladder using NF-κB p65 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.