

NIFK RABBIT PAB

货号: S218628

产品全名: NIFK 兔多抗

基因符号: Nopp34; MKI67IP

UNIPROT ID: Q9BYG3 (Gene Accession - BC022990)

背景: This gene encodes a protein that interacts with the forkhead-associated domain of the Ki-67 antigen. The encoded protein may bind RNA and may play a role in mitosis and cell cycle progression. Multiple pseudogenes exist on chromosomes 5, 10, 12, 15, and 19.

抗原: Fusion protein of human NIFK

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 200-1000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Cancer


储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human esophagus cancer tissue using 218628(NIFK Antibody) at a dilution of 1/20(Nucleus).



In comparison with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with the fusion protein and then with 218628(Anti-NIFK Antibody) at dilution 1/20.




The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using 218628(Anti-NIFK Antibody) at a dilution of 1/20.



In comparison with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with fusion protein and then with D224828(Anti-NIFK Antibody) at dilution 1/20.

kDa



Gel: 8%SDS-PAGE, Lysate: 40 µg;
Lane: A172 cell lysate;
Primary antibody: 218628(NIFK Antibody) at dilution 1/250;
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
Exposure time: 1 minute



Product Description

Pioneering GTPase and Oncogene Product Development since 2010
