

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

NIPSNAP2 RABBIT PAB

货号: S215762

产品全名: NIPSNAP2 兔多抗

基因符号 GBAS

UNIPROT ID: O75323 (Gene Accession - NP_001474)

背景: This gene encodes a member of the NipSnap family of proteins that may be involved in vesicular transport. The encoded protein is localized to mitochondria and plays a role in oxidative phosphorylation. A pseudogene of this gene is located on the long arm of chromosome 2. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

抗原: Synthetic peptide of human NIPSNAP2

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 30-150;WB: 200-1000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human cervical cancer tissue using 215762(NIPSNAP2 Antibody) at a dilution of 1/30(Cytoplasm).




In comparision with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with the synthetic peptide and then with 215762(Anti-NIPSNAP2 Antibody) at dilution 1/30



The image on the left is immunohistochemistry of paraffinembedded Human esophagus cancer tissue using 215762(Anti-NIPSNAP2 Antibody) at a dilution of 1/30.



In comparision with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with synthetic peptide and then with D163857(Anti-NIPSNAP2 Antibody) at dilution 1/30



Gel: 12%SDS-PAGE, Lysate: 40 µg; Lane 1-3: Mouse liver tissue, Human fetal liver tissue, HEPG2 cell lysates;

lysates; Primary antibody: 215762(NIPSNAP2 Antibody) at dilution 1/300; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 1 second



Product Description

Pioneering GTPase and Oncogene Product Development since 2010