

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

NPM3 RABBIT PAB

货号: S211395

产品全名: NPM3 兔多抗 基因符号 PORMIN; TMEM123

UNIPROT ID: 075607 (Gene Accession - BC054868)

背景: The protein encoded by this gene is related to the nuclear chaperone phosphoproteins, nucleoplasmin and nucleophosmin. This protein is strongly expressed in diverse cell types where it localizes primarily to the nucleus. Based on its similarity to nucleoplasmin and nucleophosmin, this protein likely functions as a molecular chaperone in the cell nucleus.

抗原: Fusion protein of human NPM3 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 100-300;WB: 500-2000;ELISA: 2000-5000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse


成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Signal Transduction, Epigenetics and Nuclear Signaling 储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human prostate cancer tissue using 211395(NPM3 Antibody) at a dilution of 1/60(Cytoplasm or Nucleus).




In comparision with the IHC on the left, the same paraffin-embedded Human prostate cancer tissue is first treated with the fusion protein and then with 211395(Anti-NPM3 Antibody) at allution 1/60.



The image on the left is immunohistochemistry of paraffinembedded Human liver cancer tissue using 211395(Anti-NPM3 Antibody) at a dilution of 1/60.



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with fusion protein and then with D122780(Anti-NPM3 Antibody) at dilution 1/60.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-2: Hela cells, Jurkat cells; Primary antibody: 211395(NPM3 Antibody) at dilution 1/300; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 10 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010