

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

P2RY1 RABBIT PAB

货号: S219959 产品全名: P2PV1 f

产品全名: P2RYI 兔多抗

基因符号 P2Y1

UNIPROT ID: P47900 (Gene Accession - NP_002554)

背景: The product of this gene belongs to the family of G-protein coupled receptors. This family has several receptor subtypes with different pharmacological selectivity, which overlaps in some cases, for various adenosine and uridine nucleotides. This receptor functions as a receptor for extracellular ATP and ADP. In platelets binding to ADP leads to mobilization of intracellular calcium ions via activation of phospholipase C, a change in platelet shape, and probably to platelet aggregation.

抗原: Synthetic peptide of human P2RY1

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-200;WB: 200-1000;ELISA: 1000-2000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction, Neuroscience

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human breast cancer tissue using 219959(P2RY1 Antibody) at a dilution of 1/40(Cytoplasm or Nucleus).




In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with the synthetic peptide and then with 219959(Anti-P2RY1 Antibody) at dilution 1/40.



The image on the left is immunohistochemistry of paraffinembedded Human colon cancer tissue using 219959(Anti-P2RYI Antibody) at a dilution of 1/40.



In comparision with the IHC on the left, the same paraffin-embedded Human colon cancer tissue is first treated with synthetic peptide and then with D260682(Anti-P2RYI Antibody) at dilution 1/40.



Gel: 10%SDS-PAGE, Lysate: 40 µg; Lane 1-2: A549 cells, mouse liver tissue; Primary antibody: 219959(P2RYI Antibody) at dilution 1/200; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 30 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010