

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010


## **PAWR RABBIT PAB**

货号: S213706 产品全名: PAWR 兔多抗 基因符号 PAR4, Par-4 UNIPROT ID: Q96IZ0 (Gene Accession - NP\_002574) 背景: The tumor suppressor WTI represses and activates transcription. The protein encoded by this gene is a WTI-interacting protein that itself functions as a transcriptional repressor. It contains a putative leucine zipper domain which interacts with the zinc finger DNA binding domain of WTI. This protein is specifically upregulated during apoptosis of prostate cells. 抗原: Synthetic peptide of human PAWR 经过测试的应用: ELISA, WB, IHC 推荐稀释比: IHC: 25-100;WB: 1000-3000;ELISA: 1000-5000 种属反应性: Rabbit 克隆性: Rabbit Polyclonal 亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human 成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol 研究领域: Epigenetics and Nuclear Signaling, Cancer 储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




cancer tissue using 213706(PAWR Antibody) at a dilution of 1/60(Cytoplasm).




The image on the left is immunohistochemistry of paraffinembedded Human ovarian cancer tissue using 213706(Anti-PAWR Antibody) at a dilution of 1/60.



Immunohistochemistry analysis of paraffin embedded Human colon In comparision with the IHC on the left, the same paraffin-embedded Human colon cancer tissue is first treated with the synthetic peptide and then with 213706(Anti-PAWR Antibody) at dilution 1/60.



In comparision with the IHC on the left, the same paraffin-embedded Human ovarian cancer tissue is first treated with synthetic peptide and then with D160694(Anti-PAWR Antibody) at dilution 1/60.



Gel: 10%SDS-PAGE, Lysate: 30 µg; Lane 1-2: Hela cells, A549 cells; Primary antibody: 213706(PAWR Antibody) at dilution 1/1200; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 10 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010